

# Neutering Your Dog

## Making an informed decision



Scroll down to read



# Neutering Your Dog – Making an Informed Decision

Dr Mark Elliott BVSc VetMFFHom MRCVS MLIHM PCH DSH RSHom

In the UK many vets are now neutering dogs routinely before their first season. As a practice we refuse to do this and to answer why, it is useful to put some facts on record.

But quite aside from these facts, our clinical experience suggests we see many more problems in neutered animals than those left entire, and even more in dogs that go through this procedure before puberty. As we see a lot of second opinion cases perhaps our views are coloured by that, but it remains a fact that neutering is banned, other than for clinical reasons, in at least 3 EU countries. It is quite arguably a mutilation with no arguable clinical benefit for the patient.

The main reason neutering is actually done, is for the perceived social benefit for the human population as:

1. It avoids some inconvenience and expense for owners.
2. It stops heat [in season] cycles.
3. No accidental pregnancies.
4. Prevents unwanted puppies being born. Some studies (USA) have shown as many as 56 per cent of litters born are unplanned;
5. It reduces the risk of abandonment of unwanted pets, resulting in mass euthanasia and neglect (but human nature will ever mean this exists somewhat as an issue), but do consider that some charities are now importing dogs from overseas to rehome here - presumably as we don't have enough unwanted dogs needing new homes in mainland Britain already?!
6. Male dogs are thought to be less likely to roam and are more likely to be better behaved - but more about that later.
7. Early neutering is said to offer breeders a chance to preclude others from using their bloodlines, allegedly also preventing use of substandard stock.
8. Vets also find that early neutering is often easier to carry out.

Point 7 is perhaps currently especially relevant as we have been seeing puppies sent to their new homes already neutered at only 8 weeks old!!! Some of these pups are proving aggressive, untrainable, allergic and suffering growth problems. It seems a problem particularly affecting some of the designer Poodle cross bred dogs and really, in our opinion, it is animal cruelty to subject these dogs to a lifetime of problems.

Since writing a former version of this article we have become increasingly concerned over the continued marketing of neutering as a health benefit despite articles in the veterinary literature debunking this myth as well as now survey after survey finding ever more problems. We would argue for decisions to be made on an individual evidence-based assessment of each patient and fully informed client consent.

The recently updated Animal Welfare Act puts a greater obligation on owners to consider their animals' health and welfare, including their ability to display normal behaviour. Clearly, certain behaviour patterns are prevented and there is no way all owners could manage their pet if all dogs were left entire, but we do need to consider some of the broad assumptions and statements about

neutering from a Canine perspective that all owners should be made aware of when deciding if neutering is best for their pet.

So does neutering benefit a dog's health, and does early neutering (before the first season in bitches and from as early as nine weeks old) offer significant benefits over a later operation or, in fact, does it cause harm as we think it does? Here are the facts I have collected together.

Please note that all figures given in this article relate to the studies I could find on the subjects (and there aren't that many), and it is always difficult to research a negative result - and even harder to get funding for such! Accordingly all figures can only be viewed as approximate and will be defined better, I hope, by future research.

The following chart is just one summary of the benefits and potential adverse events of Neutering in large- and giant-breed dogs published in the veterinary literature. Of note, much of the literature on this topic is retrospective and based on smaller select populations, so relative risk is difficult to determine for individual animals.

## **EFFECTS OF NEUTERING ON RELATIVE RISK FOR LARGE BREED DOGS**

<b>Condition</b>	<b>Effect of OHE on Relative Risk</b>	<b>Effect of Castration on Relative Risk</b>
Overall longevity	Mild increase in longevity	Mild increase in longevity
Obesity	Moderate increase	Moderate increase
Cranial cruciate ligament disease	Moderate increase*	Moderate increase*
Hip dysplasia	Mild increase*	Mild increase*
Mammary tumors	Marked decrease*	N/A
Uterine, ovarian, vaginal tumors	Prevents	N/A
Testicular tumors	N/A	Prevents
Perianal gland tumors	N/A	Marked decrease
Prostatic carcinoma	N/A	Mild increase
Lymphoma	Mild increase	Mild increase*
Mast cell tumors	Mild increase	N/A
Hemangiosarcoma	Mild increase*	Mild increase
Osteosarcoma	Mild increase*	Mild increase*
Transitional cell carcinoma	Mild increase	Mild increase
Urinary sphincter mechanism incompetence	Moderate increase*	N/A
Cystitis	Mild increase*	N/A
Benign prostatic hyperplasia	N/A	Marked decrease
Perineal hernia	N/A	Moderate decrease

\*Age at time of surgery may be important.  
 OHE stands for Ovario-Hysterectomy  
 See Further Reading D

## Is Neutering is a relatively safe procedure?

Various studies incorporating neutering show that post-operative complications following elective surgery vary considerably. Some show frequencies between 1 and 24 per cent for all complications and 1 to 4 per cent for severe complications.

Some dogs do die, but as best I could tell this is reported at around 0.1 per cent (or one in 1,000). I could find no studies showing reduced mortality risk with early neutering but as you can imagine it is very difficult to design a study to research a negative result.

## Pyometra

Ovariohysterectomy (OHE) prevents pyometra (a potentially fatal womb infection) since the uterus and ovaries have been removed. The incidence of pyometra by 10 years of age has been shown to be around 23/24 per cent which is significant. However, most cases of pyometra will be resolved by the bitch having a hysterectomy at the time, with death as a result of the condition being around 4 per cent. So, relatively speaking, whilst the risk of death from spaying routinely is 0.1 per cent the overall risk of death when older from Pyometra if left entire is possibly 1 per cent.

## Incontinence (Females – urinary sphincter mechanism incompetence (USMI))

Studies vary considerably but some report between 12 and 20 per cent of bitches are said to become incontinent to varying degrees after spaying, usually around two to three years later. Others report a much smaller incidence. Most will respond to long-term treatment, but not all. Incontinence is devastating, particularly for owners where the pet lives in close proximity and when she fails to respond to treatment it can be a reason for euthanasia or rehoming.

In general, large dogs (>15 kg) have a significantly greater risk for developing USMI than smaller dogs.<sup>4,5</sup> Although dogs that have OHE before 3 months of age show an increased risk for USMI as compared with dogs that have OHE between 3 and 12 months of age,<sup>6</sup> other data and analyses have not supported a causal link between age at time of OHE and risk for USMI.<sup>4,5,7,8</sup>

## Cystitis

Prepubescent OHE can result in a recessed or hypoplastic vulva in some dogs and may predispose these animals to perivulvar dermatitis and cystitis, particularly if they are overweight and have USMI. These findings may explain the greater reported incidence of cystitis in dogs undergoing OHE before 5.5 months of age.<sup>6</sup>

## Prostate problems

Benign prostatic hyperplasia is seen in 50% of intact males by 5 years of age<sup>9</sup>, but this does not mean it causes a concern or any problem at all. Castration prevents benign prostatic hyperplasia as well as other associated diseases (eg, prostatitis, prostatic cysts, perineal herniation).<sup>10-12</sup>

In older intact males with simple enlargement of the prostate, sometimes due to testicular tumours, late castration is usually curative, as it is with most testicular cancer. However, prostatic cancer, which logically you would think not a problem in neutered dogs, some studies suggest is actually is up to 8

times more likely (see below)!

Overall Prostatic Cancer incidence is around 0.6% negating the benefit of the castration argument when it comes to prostate issues.

### **Musculoskeletal Considerations**

Removing hormonal influence on the developing skeleton via Neutering can result in delayed physal (growth plate) closure<sup>13,14</sup> and longer-limbed conformation. The latter may play a role in the development of orthopaedic disease, as shown in studies of Labradors and Golden Retrievers neutered <6 months of age. These dogs had a 2× to 5× increased incidence of ≥1 joint disorders as compared with intact dogs.<sup>15,16</sup>

### **Canine Cranial Cruciate Ligament Disease**

Large-breed dogs that underwent Neutering at <6 months of age have shown a 3× increased risk for excessive tibial plateau angle and predisposition for earlier canine cranial cruciate ligament (CCL) injury.<sup>17</sup> Dogs that underwent Neutering at a non-specified age had a 2× to 3× incidence of CCL disease as compared with intact dogs.<sup>18,19</sup> In a study of 750 Golden Retrievers, none of the intact dogs had CCL disease, compared with an incidence of 5% in castrated dogs and 7.7% in spayed dogs that underwent neutering at <12 months of age.<sup>16</sup> Body Condition Score (BCS) was the same for dogs with and without CCL disease. This suggests that change in conformation—not just increased body weight associated with neutering—was responsible.<sup>16</sup>

### **Hip Dysplasia**

Hip dysplasia may be influenced by patient sex and breed as well as timing of neutering. In the Golden Retriever study,<sup>16</sup> incidence of hip dysplasia in males neutered at <12 months of age was double that of intact males, with an earlier onset of disease. The BCS of the males with and without hip dysplasia and neutered at <12 months of age was not far greater. No significant difference in hip dysplasia incidence was seen in the females.<sup>16</sup> Spayed or neutered Boxers with a mean age of 3 years at the time of neutering had a 1.5× increased risk for developing hip dysplasia.<sup>20</sup> Data collected from a Veterinary Medical Database between 1964 and 2003 showed that neutering (at a non-specified age) increased the likelihood of hip dysplasia by 17%.<sup>18</sup> Incidence of hip dysplasia was 6.7% in dogs that underwent neutering before 5.5 months of age and 4.7% in dogs that underwent neutering between 5.5 months and 1 year of age.<sup>6</sup>

### **Obesity**

Obesity plays a significant role in the development and progression of many orthopaedic diseases and osteoarthritis.<sup>21</sup> Although neutering is a significant risk factor for obesity,<sup>6,22,23</sup> neutering alone is most likely less important than other environmental factors (eg, diet, exercise regimen).<sup>24</sup>

## **Cancers**

### **Mammary Tumors**

Many Vets are aware of the effect and timing of spaying on incidence of mammary tumors based on Schneider, Dorn, and Taylor's 1969 study.<sup>25</sup> This is argued so often as a significant reason for neutering but does it really stack up?

It is said that spaying after the third estrous cycle and after 2.5 years of age appears to provide minimal protection against mammary tumour development.<sup>25,26</sup> However, a 2010 systematic review of this and other studies on the protective effect of OHE concluded that the evidence is weak because of confounding factors and bias.<sup>27</sup>

Whatever, the statements are somewhat misleading. Unneutered bitches have only a 3.4 per cent chance of developing this problem with age, with 50 per cent of those cases being malignant. Early neutering reduces this risk to 0.5 per cent of the 3.4 per cent (= 0.017 per cent), but spaying the bitch after the first season and before the second reduces it also to eight per cent of 3.4 per cent (= 0.27 per cent).

So the argument for early neutering over leaving the bitch to have one season on the basis of this without considering all the other issues doesn't really stack up as particularly significant to my mind.

There is also a downside as regards to Cancer generally which is that some studies show increased incidence of other cancers following neutering including some of those discussed below.

### **Reproductive Tumors & Tumors Influenced by Hormones**

Neutering eliminates the potential for developing uterine, ovarian, and testicular tumors through removal of the primary organ.<sup>31</sup> Perianal gland tumors in male dogs are treated successfully via castration.<sup>32</sup> OHE is protective against vaginal leiomyomas and can decrease recurrence, even with incomplete surgical resection.<sup>33,34</sup> Neutered male dogs had 2× to 8× the incidence of prostatic carcinoma as compared with intact male dogs<sup>34,35</sup>; however, the overall prevalence of prostatic cancer is <1%.<sup>35-37</sup>

### **Lymphoma**

A large population study showed that intact female dogs had a significantly lower risk for developing lymphoma as compared with dogs that underwent Neutering (at a nonspecific age) or intact male dogs.<sup>38</sup> This finding was consistent in studies of golden retrievers and vizslas, although castration at <12 months of age was also found to be a risk factor.<sup>16,39</sup>

### **Mast Cell Tumors**

Neutering has been associated with 2× to 4× the risk for mast cell tumors, particularly in female dogs.<sup>16,39,40</sup> However, estrogen receptors have not been identified in mast cell tumors, so a direct hormonal link has not been established.<sup>41</sup>

### **Heamangiosarcoma**

Golden retrievers that underwent OHE after 1 year of age had 4× the incidence of hemangiosarcoma as compared with intact females or females that underwent OHE before 1 year of age.<sup>16</sup> No significant differences in incidence of hemangiosarcoma were found in male Golden Retrievers.<sup>16</sup> Similar findings

were noted in a study of Vizslas, although dogs that underwent OHE before 1 year of age or castration after 1 year of age also had increased risk.<sup>39</sup> Other non-breed-specific studies have shown similar findings for splenic and cardiac hemangiosarcoma.<sup>42,43</sup>

### **Osteosarcoma**

An increased risk for osteosarcoma was seen in Rottweilers that underwent OHE or castration before 1 year of age, although the overall 13% incidence of bone sarcomas in this study group seems disproportionately high.<sup>44</sup> Historic studies have reported a 1.3× to 1.9× increased risk for osteosarcoma in animals that underwent neutering at a non-specified age.<sup>45,46</sup>

### **Transitional Cell Carcinoma**

Female dogs are more predisposed to bladder transitional cell carcinoma than male dogs, and neutering (at a non-specified age) increases the risk up to 3× in both male and female dogs.<sup>47,48</sup> An 8× increase in prostatic transitional cell carcinoma has also been reported in male dogs that underwent castration at a non-specified age.<sup>36</sup>

### **Neutered pets are better behaved, and early neutered ones better still – NOT SO!**

Quite apart from the fact that most behaviour problems are created by owners failing to understand and/or train their dogs properly, knowing factors relative to the breed and individual characteristics, and possibly also failure to provide an appropriate environment, there are a number of arguments to consider here.

Yes, neutered male dogs are less likely to take it upon themselves to roam freely and, yes, they are less likely to exhibit normal behaviour and try to assert male dominant behaviour, but all these things can be controlled by human interventions.

However, the American Kennel Club Canine Health Foundation reported significantly more behavioural problems in neutered dogs and bitches. Other studies have also shown early neutering to be associated with increased incidence of noise phobias and undesirable sexual behaviour! Quite the opposite of what one would logically expect if neutering was a benefit. The most commonly observed problem in spayed bitches was found to be fearful behaviour, and in male dogs aggression (the very thing we castrate them for!). I wonder if some of the latter could well be due to confusion as to the individual's place in dog society, and for early neutered dogs the existence as an adult-sized animal locked into a state of permanent puppyhood?

On the upside, separation anxiety and inappropriate toileting when frightened have been found to improve. Perhaps the result of a suppression of mental development?

### **Ageing**

Longer term, studies have also shown sexually intact males to show slower cognitive impairment with age than neutered dogs. Results with bitches were inconclusive.

## Hypothyroidism

Studies on several thousand Golden Retrievers showed that those neutered were more likely to develop hypothyroidism and this affects mental function amongst its other concerns. Certainly I have diagnosed early neutered dogs that have developed this condition as early as seven months of age, and also have seen early neutered dogs fail expensive training programmes when they develop it young. This perhaps is the reason why behaviourists are telling me they are being called upon to help owners finding these early neutered individuals so hard to train?

Other studies have confirmed this **Hypothyroid** link and, for interest, the most common clinical findings in hypothyroidism include obesity, seborrhoea (greasy skin), alopecia (hair loss), weakness, lethargy, bradycardia (slow heart rate), and pyoderma (skin infection). Skin diseases make up a lot of the work in current veterinary practice and most dogs are neutered – as they say in the USA “go figure”.

## Dachshunds

These little dogs do sadly seem to suffer hugely as a result of neutering, beyond that quoted for some of the larger breeds

Intervertebral Disc Disease (IVDD) is the key issue affecting Dachshunds, confirmed by the 2014 Kennel Club Survey (E). Dachshunds are relatively long lived, with the oldest standard being 16 years old, and the oldest miniature being 19 on survey. Sadly dogs that suffer IVDD are typically euthanased in the 4-6 year age range (E) When you consider that 17% of Dachshunds have some degree of back issue and the odds of a neutered dog over the age of 3 of suffering this condition is nearly double (1.8x) a lot of dogs are dying young as a result of being spayed or castrated (F)

What is perhaps truly shocking is the effect of early neutering. Further analysis of data from 2015 was reported in the 2016 Dachshund Breed Council Health Report and this showed that when the 4 year old dogs in the survey were looked at those neutered under 6 months of age were 12 times more likely to have suffered an IVDD incident than entire dogs. Looking at the 4-5 year olds 25% of early neutered dogs have suffered IVDD compared to 7% of entire Dachshunds.

The 2016 report went on to analyze other data as well. This showed NO statistically significant difference in mammary tumour incidence between spayed and entire bitches, although the numbers show spayed bitches were more likely to have the disease. Neutering also increased the odds of a Dachshund having skin allergies and autoimmune conditions by factors of 2x and 4x.

## Autoimmune Disorders

A recent paper ( ) analyzing patient records from 90,090 dogs from a teaching hospital in California over a 15 year period looked at the prevalence and risk of atopic dermatitis (ATOP), autoimmune hemolytic anemia (AIHA), canine myasthenia gravis (CMG), colitis (COL), hypoadrenocorticism (ADD), hypothyroidism (HYPO), immune-mediated polyarthritis (IMPA), immune-mediated thrombocytopenia (ITP), inflammatory bowel disease (IBD), lupus erythematosus (LUP), and pemphigus complex (PEMC), for intact females, intact males, neutered females, and neutered males.

Neutered dogs had a significantly greater risk of ATOP, AIHA, ADD, HYPO, ITP, and IBD than intact

dogs with neutered females being at greater risk than neutered males for all but AIHA and ADD. Neutered females, but not males, had a significantly greater risk of LUP than intact females.

That study concluded neutering is associated with increased risk for certain autoimmune disorders and underscore the need for owners to consult with their veterinary practitioner prior to neutering to evaluate possible benefits and risks associated with such a procedure.

Dare I rest my case here?

I could go on, but it would be impossible within the scope of this article to cover all the possible other minor issues seen as a result of neutering, so I shall stop at this point! Readers can easily find all of this information and more on the Internet.

### **In Conclusion**

It is important that owners make an informed choice for their pet, but I would venture to ask 'where are the benefits for the individual dog in all this?' It should be accepted that most of the benefits are clearly associated with societal, human behaviour and convenience arguments. Many years ago dogs were domesticated and perhaps this is the price they pay as part of that bargain?

Clinically the arguments for neutering dogs do not, in my opinion, stack up to much at all and it may actually be a negative action when looking at their long term health. While for bitches there are some apparent positive benefits, they are still minimal in my opinion, when compared to the risk factors and long term health issues should the individual be unfortunate enough to suffer them.

Ultimately the choice an owner makes (and the choice should be the owners) will mostly depend on personal circumstances, the breed and how you wish the dog to interact with your family – all human factors.

With regards to early neutering owners should be aware of the long term issues that appear to be associated with early neutering over letting the dog mature, and be prepared to cope should they happen. Promotion of early neutering without explanation of the risks is in my opinion unethical, and we have decided as a practice we will not do it anyway as the negatives are just too many and so it is contrary to our ethos.

A decision as regards to whether or not to have your dog neutered should be an informed one - it does not have to follow others' agendas and media campaigns - and, hopefully, I have gone some way to helping with that.

## REFERENCES

- 1 Trevejo R, Yang M, Lund EM. Epidemiology of surgical castration of dogs and cats in the United States. *JAVMA*. 2011;238(7):898-904.
- 2 Spain CV, Scarlett JM, Cully SM. When to neuter dogs and cats: a survey of New York state veterinarians' practices and beliefs. *JAAHA*. 2002;38(5):482-488.
- 3 Olson PN, Kustritz MV, Johnston SD. Early-age neutering of dogs and cats in the United States: a review. *J Reprod Fertil Suppl*. 2001;57:223-232.
- 4 Forsee KM, Davis GJ, Mouat EE, Salmeri KR, Bastian RP. Evaluation of the prevalence of urinary incontinence in spayed female dogs: 566 cases (2003-2008). *JAVMA*. 2013;242(7):959-962.
- 5 Stöcklin-Gautschi NM, Hässig M, Reichler IM, Hubler M, Arnold S. The relationship of urinary incontinence to early spaying in bitches. *J Reprod Fertil Suppl*. 2001;57:233-236.
- 6 Spain CV, Scarlett JM, Houpt KA. Long-term risks and benefits of early-age Neutering in dogs. *JAVMA*. 2004;224(3):380-387.
- 7 Beauvais W, Cardwell JM, Brodbelt DC. The effect of neutering on the risk of urinary incontinence in bitches: a systematic review. *J Small Anim Pract*. 2012;53(4):198-204.
- 8 de Bleser B, Brodbelt DC, Gregory NG, Martinez TA. The association between acquired urinary sphincter mechanism incompetence in bitches and early spaying: a case-control study. *Vet J*. 2011;187(1):42-47.
- 9 Berry SJ, Strandberg JD, Saunders WJ, Coffey DS. Development of canine benign prostatic hyperplasia with age. *Prostate*. 1986;9(4):363-373.
- 10 Nizański W, Levy X, Ochota M, Pasikowska J. Pharmacological treatment for common prostatic conditions in dogs—benign prostatic hyperplasia and prostatitis: an update. *Reprod Domest Anim*. 2014;49Suppl2:8-15.
- 11 Mann FA, Nonneman DJ, Pope ER, Boothe HW, Welshons WV, Ganjam VK. Androgen receptors in the pelvic diaphragm muscles of dogs with and without perineal hernia. *Am J Vet Res*. 1995;56(1):134-139.
- 12 Snell WL, Orsher RJ, Larenza-Menzies MP, Popovitch CA. Comparison of caudal and pre-scrotal castration for management of perineal hernia in dogs between 2004 and 2014. *N Z Vet J*. 2015;63(5):272-275.
- 13 Root MV, Johnston SD, Olson PN. The effect of prepuberal and postpuberal Neutering on radial physeal closure in male and female domestic cats. *Veterinary Radiology & Ultrasound*. 1997;38(1):42-47.
- 14 Salmeri KR, Bloomberg MS, Scruggs SL, Shille V. Neutering in immature dogs: effects on skeletal, physical, and behavioral development. *JAVMA*. 1991;198(7):1193-1203.
- 15 Hart BL, Hart LA, Thigpen AP, Willits NH. Long-term health effects of neutering dogs: comparison of Labrador retrievers with golden retrievers. *PLoS One*. 2014;9(7):e10224.
- 16 Torres de la Riva G, Hart BL, Farver TB, et al. Neutering dogs: effects on joint disorders and cancers in golden retrievers. *PLoS One*. 2013;8(2):e55937.
- 17 Duerr FM, Duncan CG, Savicky RS, Park RD, Egger EL, Palmer RH. Risk factors for excessive tibial plateau angle in large-breed dogs with cranial cruciate ligament disease. *JAVMA*. 2007;231(11):1688-1691.
- 18 Witsberger TH, Villamil JA, Schultz LG, Hahn AW, Cook JL. Prevalence of and risk factors for hip dysplasia and cranial cruciate ligament deficiency in dogs. *JAVMA*. 2008;232(12):1818-1824.
- 19 Slauterbeck JR, Pankratz K, Xu KT, Bozeman SC, Hardy DM. Canine ovariohysterectomy and orchiectomy increases the prevalence of ACL injury. *Clin Orthop Relat Res*. 2004;(429):301-305.
- 20 van Hagen MA, Ducro BJ, van den Broek J, Know BW. Incidence, risk factors, and heritability estimates of hind limb lameness caused by hip dysplasia in a birth cohort of boxers. *Am J Vet Res*. 2005;66(2):307-312.

- 21 Frye CW, Shmalberg JW, Wakshlag JJ. Obesity, exercise and orthopedic disease. *Vet Clin North Am Small Anim Pract.* 2016; doi: 10.1016/j.cvsm.2016.04.006.
- 22 Lefebvre SL, Yang M, Wang M, Elliott DA, Buff PR, Lund EM. Effect of age at Neutering on the probability of dogs becoming overweight. *JAVMA.* 2013;243(2):236-243.
- 23 McGreevy PD, Thomson PC, Pride C, Fawcett A, Grassi T, Jones B. Prevalence of obesity in dogs examined by Australian veterinary practices and the risk factors involved. *Vet Rec.* 2005;156(22):695-702.
- 24 Courcier EA, Thomson RM, Mellor DJ, Yam PS. An epidemiological study of environmental factors associated with canine obesity. *J Small Anim Pract.* 2010;51(7):362-367.
- 25 Schneider R, Dorn CR, Taylor DO. Factors influencing canine mammary cancer development and postsurgical survival. *J Natl Cancer Inst.* 1969;43(6):1249-1261.
- 26 Sonnenschein EG, Glickman LT, Goldschmidt MH, McKee LJ. Body conformation, diet, and risk of breast cancer in pet dogs:a case-control study. *Am J Epidemiol.* 1991;133(7):694-703.
- 27 Beauvais W, Cardwell JM, Brodbelt DC. The effect of neutering on the risk of mammary tumours in dogs: a systematic review. *J Small Anim Pract.* 2012;53(6):314-322.
- 28 Egenvall A, Bonnett BN, Ohagen P, Olson P, Hedhammar A, von Euler H. Incidence of and survival after mammary tumors in a population of over 80 000 insured female dogs in Sweden from 1995 to 2002. *Prev Vet Med.* 2005;69(1-2):109-127.
- 29 Brønden LB, Nielsen SS, Toft N, Kristensen AT. Data from the Danish veterinary cancer registry on the occurrence and distribution of neoplasms in dogs in Denmark. *Vet Rec.* 2010;166(19):586-590.
- 30 Benjamin SA, Lee AC, Saunders WJ. Classification and behavior of canine mammary epithelial neoplasms based on life-span observations in beagles. *Vet Pathol.* 1999;36(5):423-436.
- 31 Smith AN. The role of neutering in cancer development. *Vet Clin North Am Small Anim Pract.* 2014;44(5):965-975.
- 32 Wilson GP, Hayes HM Jr. Castration for treatment of perianal gland neoplasms in the dog. *JAVMA.* 1979;174(12):1301-1303.
- 33 Thacher C, Bradley RL. Vulvar and vaginal tumors in the dog:a retrospective study. *JAVMA.* 1983;183(6):690-692.
- 34 Herron MA. Tumors of the canine genital system. *JAAHA.* 1983;19:981-994.
- 35 Teske E, Naan EC, van Dijk EM, Van Garderen E, Schalken JA. Canine prostate carcinoma: epidemiological evidence of an increased risk in castrated dogs. *Mol Cell Endocrinol.* 2002;197(1-2):251-255.
- 36 Bryan JN, Keeler MR, Henry CJ, Bryan ME, Hahn AW, Caldwell CW. A population study of neutering status as a risk factor for canine prostate cancer. *Prostate.* 2007;67(11):1174-1181.
- 37 Cornell KK, Bostwick DG, Cooley DM, et al. Clinical and pathologic aspects of spontaneous canine prostate carcinoma: a retrospective analysis of 76 cases. *Prostate.* 2000;45(2):173-183.
- 38 Villamil JA, Henry CJ, Hahn AW, et al. Hormonal and sex impact on the epidemiology of canine lymphoma. *J Cancer Epidemiol.* 2009:591753.
- 39 Zink MC, Farhooody P, Elser SE, et al. Evaluation of the risk and age of onset of cancer and behavioral disorders in gonadectomized Vizslas. *JAVMA.* 2014;244(3):309-319.
- 40 White CR, Hohenhaus AE, Kelsey J, Procter-Gray E. Cutaneous MCTs: associations with spay/neuter status, breed, body size, and phylogenetic cluster. *JAAHA.* 2011;47:210-216.
- 41 Larsen AE, Grier RL. Evaluation of canine mast cell tumors for presence of estrogen receptors. *Am J Vet Res.* 1989;50(10):1779-1780.
- 42 Prymak C, McKee LJ, Goldschmidt MH, Glickman LT. Epidemiologic, clinical pathologic, and prognostic characteristics of splenic heman giosarcoma and splenic hematoma in dogs. *JAVMA.* 1988;193:706-712.
- 43 Ware WA, Hopper DL. Cardiac tumors in dogs: 1982-1995. *JVIM.* 1999;13(2):95-103.

- 44 Cooley DM, Beranek BC, Schlittler DL, Glickman NW, Glickman LT, Waters DJ. Endogenous gonadal hormone exposure and bone sarcoma risk. *Cancer Epidemiol Biomarkers Prev.* 2002;11(11):1434-1440.
- 45 Priester WA, McKay FW. The occurrence of tumors in domestic animals. *Natl Cancer Inst Monogr.* 1980;(54):1-210.
- 46 Ru G, Terracini B, Glickman LT. Host related risk factors for canine osteosarcoma. *Vet J.* 1998;156(1):31-39.
- 47 Norris AM, Laing EJ, Valli VE, et al. Canine bladder and urethral tumors: a retrospective study of 115 cases (1980-1985). *JVIM.* 1992;6(3):145-153.
- 48 Mutsaers AJ, Widmer WR, Knapp DW. Canine transitional cell carcinoma. *JVIM.* 2003;17(2):136-144.
- 49 Hoffman JM, Creevy KE, Promislow DE. Reproductive capability is associated with lifespan and cause of death in companion dogs. *PLoS One.* 2013;8(4):e61082. Lifetime golden retriever study. Morris Animal Foundation. <http://caninelifetimehealth.org>. Updated 2016. Accessed November 2015.

#### Further Reading

- A. Long term risks and benefits of early age gonadectomy in dogs. C.V.Spain et al. *JAVMA* Vol 224, No3, Feb1, 2004
- B. Canine Reproduction. Peggy Root DVM. Article for the Puli club of America <http://www.puliclub.org/CHF/AKC2007Conf/Canine%20Reproduction.htm>
- C. Early Spay-Neuter Considerations for the Canine Athlete. C Zink 2005 <http://www.caninesports.com/Spayneuter.html>
- D. Age of Neutering in Large and Giant-Breed Dogs. Article in Clinicians Brief by Clara S.S.Goh BVSc MS DACVS, ACVS Founding Fellow (Surgical Oncology), Colorado State University. August 2016. From which many of the references and some of the text in this article are taken.
- E. Kennel Club Survey 2014 Dachshund Health
- F. DachsLife 2015: The Dachshund Breed Council Lifestyle and Health Survey (Online) and associated open access paper - DachsLife 2015: an investigation of lifestyle associations with the risk of intervertebral disc disease in Dachshunds by R. M. A. Packer, I. J. Seath, D. G. O'Neill, S. De Decker and H. A. Volk *Canine Genetics and Epidemiology* 2016 3:8 <https://doi.org/10.1186/s40575-016-0039-8>
- G. The Dachshund Breed Council Health Report 2016
- F. Gonadectomy effects on the risk of immune disorders in the dog: a retrospective study, Crystal R. Sundburg, Janelle M. Belanger, Danika L. Bannasch, Thomas R. Famula and Anita M. Oberbauer. *BMC Veterinary Research* 2016 12:278 <https://doi.org/10.1186/s12917-016-0911-5>